

Colonoscopy Bowel Preparation

Morning appointment between 10am and 12 pm



For a **successful colonoscopy** it is very important that the lining of the bowel is clearly visible so that the risk of lesions being missed is minimised. Poor preparation may mean that the procedure will have to be stopped and repeated on another day, or that the accuracy of colonoscopy is reduced.

Bowel preparation – where to get your bowel prep?

Option 1	The bowel preparation can be dispensed free of charge from the Redcliffe Hospital Pharmacy. You may need to wait 30-60 minutes.	To access these videos, you can scan the QR code below with your smart device or alternatively you can type https://metronorth.health.qld.gov.au/redcliffe/healthcare-services/gastroenterology into your internet search engine on a computer or smart device. It is important that you follow the instructions given by the Redcliffe Hospital Endoscopy department as your circumstances may be different from the general information supplied in these videos. 
Option 2	The bowel preparation can be purchased at your own cost from your local pharmacy using Colonoscopy Preparation Order Guide on the last page of these instructions. • Please be advised we are unable to send medication in the post. • Disregard the instructions on the Glycoprep-O kit and follow these instructions only.	
Watch a video to help you prepare for your Colonoscopy. To help you prepare for your colonoscopy our medical teams and patients have created a series of films about colonoscopies. Each film has been reviewed by clinicians and patients alike to ensure that the information shared can help you to understand the procedure and prepare for it.		

7 Days prior to your appointment

- Start taking Movicol (if advised by a clinician) this will help to prepare your bowel prior to commencing your bowel preparation. Take one sachet in the morning and one sachet at night.
- You must cease taking Iron tablets, herbal supplements, multivitamins, and Fish/Krill Oil.
- Please continue to take your regular medications as prescribed unless otherwise advised by a clinician.

Important information – contact Redcliffe Hospital Endoscopy Department on 3883 7800

- If you are taking any blood thinning medication other than aspirin, and you have not already received instructions prior to your appointment. **DO NOT stop taking anticoagulant medication unless instructed to by a clinician.**
- If you are taking diabetic tablets or insulin, and you have not already received instructions on diabetes management prior to colonoscopy.
- If you are pregnant or breastfeeding.

Tips for drinking Bowel Preparation

- Keep yourself warm whilst drinking the bowel preparation.
- Drink the solution through a straw, suck the occasional barley sugar or clear peppermint between drinks.
- Apply protective cream, e.g. Lanoline around your anal area to reduce excoriation.
- If you feel nauseated whilst taking the preparation stop for ½ hour. Have a hot drink & walk around, resume drinking & continue as before. If you anticipate that you may get nauseous while doing preparation, please see your GP to

have anti-nausea medication ready prior to commencing your preparation.

- Visit your GP or pharmacy if you have trouble tolerating your bowel preparation.
 - If you feel bloated, try walking around, suck some clear peppermints / drink peppermint tea.
 - If you become ill & have severe pain seek medical advice from your GP or nearest Emergency Medical Centre.
- If you have haemorrhoids (piles) use your haemorrhoid cream to reduce discomfort.

4 days before your procedure – Start a low fibre diet – **Avoid all foods with nuts, seeds and skin.**

Bread & Cereals – Choose ✓ Plain white bread / toast ✓ Plain muffins ✓ Plain bread rolls ✓ Crumpets ✓ Plain croissants ✓ Pita bread ✓ Refine breakfast cereals, Semolina, Cornflakes, Rice bubbles etc. ✓ Plain cakes and biscuits ✓ Plain pancakes / pikelets ✓ White rice ✓ White pasta ✓ Couscous ✓ Polenta ✓ Bagels Bread & Cereals -Avoid ✗ Wholemeal, multigrain, brown and Rye bread ✗ Wholegrain cereals e.g. All-Bran, Weetbix etc ✗ Cakes, muffins and biscuits with Nuts or dried fruit ✗ Brown rice or pasta ✗ Unprocessed bran / rolled oats ✗ Porridge or muesli ✗ Brown crumpets ✗ Wholemeal flour or oatmeal	Fruit and Vegetables – Choose ✓ Potato / pumpkin without skin ✓ Marrow / squash / pawpaw ✓ Choko ✓ Avocado ✓ Mushrooms ✓ Ripe banana ✓ Peaches, pears and apples without skin ✓ Rockmelon / watermelon ✓ Canned peaches, apricots, apples and pears Fruit & Vegetables – Avoid ✗ All other fruit and vegetables, including salad vegetables and dried fruit Drinks - Choose ✓ Water ✓ Tea & Coffee ✓ Milk drinks, soft drinks ✓ Cordial, Strained juices ✓ Clear soups ✓ Milo, Ovaltine Drinks – Avoid ✗ Unstrained fruit juices ✗ Red and green coloured drinks	Meat & other proteins – Choose ✓ Beef, lamb, veal, pork, poultry, eggs, fish and shellfish Meat & other proteins – Avoid ✗ Baked beans lentils and other legumes ✗ Casseroles or rissoles ✗ Stir fry meals ✗ Nuts & seeds ✗ Pizza Dairy Foods – Choose ✓ Milk, cheese, plain yoghurt ✓ Ice cream, custard ✓ Cream, butter / margarine ✓ Condensed/evaporated milk Dairy Foods – Avoid ✗ Dairy products with fruit, nuts or seeds Other Foods – Choose ✓ Plain/clear lollies (not red) ✓ Sugar, honey or chocolate ✓ Strained jams/jelly ✓ Oils and spices Other foods – Avoid ✗ Peanut butter, jams with skins and seeds or marmalade
--	--	---



The day before your procedure – Clear Fluids all day – NO SOLID FOOD

- You cannot have any solid foods. A clear fluid ONLY diet must be followed.
- **Stop taking the Movicol if you were required to take it.**
- Prepare your bowel preparation solution in the morning and refrigerate until ready to drink.
 - Dissolve the **Magnesium citrate sachet** in **1 glass of hot water (250mL).**
 - Mix the three sachets of **GLYCOPREP-ORANGE** in **3 litres of water.**
- It is important to drink adequate fluids to avoid headaches /dehydration. You may drink as much clear fluids as you like throughout the day including when drinking bowel preparation. Examples of clear fluids:

- ✓ Clear strained soup
- ✓ Herbal tea or iced tea
- ✓ Jelly (green or yellow only)
- ✓ Lemonade or ginger ale

- ✓ Water or clear apple juice
- ✓ Yellow or orange sports drinks
- ✓ Black tea or black coffee
- ✓ Orange or lemon cordial.

3 PM - Drink the **Magnesium citrate** and take the **3 yellow Bisacodyl tablets** with 1 full glass of water.



5 PM - Drink **2 Litres of GLYCOPREP-ORANGE** (reserve the 3rd litre). This will produce watery diarrhoea. Drink approximately 1 glass every 15 minutes. You should complete this by 7:30 pm. If having trouble tolerating bowel preparation, you can pause until the symptoms settle, then restart.



From the time you finish drinking the solution and until you go to bed you should drink at least 3 glasses of water or other clear fluids to prevent dehydration. Diarrhoea should settle around 9.30 PM.

Day of Procedure


You must remain on a clear fluid diet

X NO JELLY, BARLEY SUGARS OR CLEAR PEPPERMINTS DAY OF PROCEDURE

- ✓ Herbal tea or iced tea
- ✓ Lemonade or ginger ale
- ✓ Water or clear apple juice

- ✓ Yellow or orange sports drinks
- ✓ Black tea or black coffee
- ✓ Orange or lemon cordial

5 AM - Drink the remaining **1 Litre of GLYCOPREP-ORANGE**, you should finish this by **6 AM**. Continue to drink **clear fluids**.



8 AM - NIL BY MOUTH – Do not drink anything. Your bowel motions should be clear (or look like urine) and be free of particles. You can continue to sip fasting fluids (maximum 170ml/hr) until your procedure.



Bowel motions NOT CLEAR enough

YOU ARE READY 😊

If your bowel motion is not clear after completing your bowel preparation please call ☎ (07) 3883 7800

Following your colonoscopy procedure

- ✓ You must have a responsible adult to pick you up and stay with you overnight.
- ✓ You will not be able to undertake the following activities for 24 hours after your procedure: Drive any vehicle, use heavy machinery, sign legal documentation or drink alcohol.
- ✓ You may resume normal eating habits after your procedure, unless otherwise instructed.

Colonoscopy Bowel Preparation – Order Guide

To purchase your bowel preparation, please take this form into a pharmacy and the pharmacist will know what bowel preparation to provide.

The instructions on the bowel preparation kit may differ to the instructions given to you by your doctor/hospital. It is important to always follow the instructions provided by your doctor/hospital. If you can't find your bowel preparation at your local pharmacy, ask your pharmacist to order it for you.

The codes your pharmacist will need are provided below and it should arrive within 24-48 hours.

If any concerns or if you would like to obtain your bowel preparation free of charge from Redcliffe Hospital, please contact Endoscopy Services ☐ (07) 3883 7800.

Product



Description

Glycoprep-O Kit®

3 x Glycoprep-O® 70g sachet. Each sachet contains: macrogol 3350 52.9g, sodium chloride 2.6g, potassium chloride 740mg, sodium sulfate 5.6g
3x Bisacodyl BP 5mg tablets
1x Magnesium citrate sachet 21.5g containing: magnesium carbonate 7.2g, citric acid 14.3g

Pharmacy Ordering Codes

API: 70851

Symbion: 946885

Sigma: 10000943

Please note: if Movicol was indicated as part of your bowel preparation this would need to be purchased from the pharmacy and commenced 7 days prior to your procedure date.

SERVICES

Gastroenterology Bookings

Day Procedure Unit

Preadmission Clinic

Redcliffe Hospital

CONTACT NUMBERS

3883 7800

3883 7841

3883 7345

3883 7777