Preparation for Colonoscopy Colon Consent Clinic - Split Prep

Why?

For a **successful colonoscopy** it is very important that the bowel be cleaned well. This will enable the doctor to clearly see the lining of the bowel.

Poor preparation may mean that the procedure will have to be stopped and you will have to repeat the procedure on another day.

What do I need?

Bowel Prep Pack containing

- **GLYCOPREP-C KIT**
 - √ 1 sachet Magnesium citrate
 - ✓ 3 Bisacodyl tablets (yellow)
 - ✓ 1 sachet GLYCOPREP-C

How?

By following a **low fibre diet** for 3 days. This reduces the fibre in your bowel, which makes it easier when you start the 2-day bowel-cleansing regime.

Cease Iron tablets/multivitamins, Fish/Krill Oil supplements 10 days prior

Follow low fibre diet Day 1-3

Day 4 Start the bowel preparation

Day of procedure Day 5

The bowel cleansing preparation causes you to have watery bowel motions over a short period of time.

When?

Commence the low fibre diet 4 days before the day of your scheduled procedure.

Day 1-3**Low Fibre Diet**

Start the low fibre diet. We have a provided a list of foods to assist you in this process. Eat the foods from the Choose box.

BREAD & CEREALS

Choose

- Plain white bread / toast
- Plain muffins
- Plain bread rolls
- Crumpets
- Plain croissants
- Pita bread
- Refined breakfast cereals; Semolina, Cornflakes, Rice **Bubbles** etc

- Plain cakes and biscuits
- ✓ Plain pancakes / pikelets
- White rice.
- White pasta
- Couscous
- Polenta
- Bagels

Avoid

- Wholemeal, brown and rye breads
- Multi-grain breads & muffins
- Wholegrain cereals, e.g. bran cereals, Weetbix. Vita brits
- cakes and biscuits containing nuts, dried fruit

- Brown rice
- Brown pasta
- Unprocessed bran, rolled oats
- Porridge
- Muesli
- Brown crumpets
- Muffins with fruit
- 8 Wholemeal flour and oatmeal.

FRUIT & VEGETABLES

You will need to remove the skins and seeds from these fruit and vegetables

Choose

- Potato without skin
- Pumpkin without skin
- marrow / squash
- choko
- avocado
- mushrooms
- ripe banana
- apple with no skin

- peach with no skin
- pears
- pawpaw
- rockmelon
- watermelon
- canned peaches, apples, apricots &
 - pears

- **Avoid**
- All other fruit and vegetables, including salad vegetables

MEAT & OTHER PROTEIN FOODS

Choose

- beef
- veal
- poultry
- lamb

fish & shellfish

Rissoles

Stir fry meals

pork

eggs

Avoid

- Baked Beans
- \odot Casseroles
- Lentils
- Nuts & seeds \odot
- Legumes

Pizza

DAIRY FOODS

Choose

- ✓ Milk
- ✓ Ice cream
- ✓ Plain or vanilla yoghurt
- ✓ Cheese processed & plain
- Cottage cheese

Avoid

Dairy products with fruit and seeds

DRINKS

Choose

- ✓ Water
- ✓ Tea
- ✓ Coffee
- ✓ Plain milk drinks
- ✓ Soft drinks

✓ Cordial

✓

✓

Custard

Rice pudding

Condensed milk

Evaporated milk

Powdered milk

Butter / margarine

Cream or sour cream

- ✓ Strained fruit juices
- ✓ Clear soups
- ✓ Milo or Ovaltine
- Soft drinks ✓ White wine

Avoid

- Unstrained fruit juices

OTHER FOODS

Choose

- ✓ Plain Iollies
- ✓ Clear Jellies not red

Jams with skins &

✓ Sugar

Avoid

√ Vegemite

Honey,

✓ Chocolate

seeds

Peanut paste/butter

Jellied strained jams

- Dried Fruits/ Nuts
- B Marmalade

Day 4 The day before your test

CLEAR FLUIDS ALL DAY

 It is important to drink adequate fluids to avoid dehydration. You may drink as much clear fluids throughout the day

Clear Fluids

- Clear or strained soup
- Jelly (green or yellow only)
- Water or strained fruit juice
- Black tea / coffee or soft drink
- Herbal tea / Iced tea
- Lemonade/ginger ale
- Yellow/orange Sports drinks
- Orange / Lemon cordial

8 AM Prepare

- The Magnesium citrate by dissolving in 1 glass of hot water (250mL) & refrigerate until ready to drink.
- 2. Make sachet of GLYCOPREP-C in 3 litres of water and refrigerate until ready to drink.
- 3 PM Drink the Magnesium citrate drink
 Take the 3 yellow Bisacodyl tablets with 1 full
 glass of water
- **5 PM** Begin drinking 2 litres of the **GLYCOPREP-C** solution at the rate of 1 litre / hour. This will produce watery diarrhoea within an hour.
 - You should have finished drinking this by 7:30 PM and the diarrhoea will settle around 9:30 PM.
 - From now until you go to bed you should drink 3 glasses of water or clear tea / coffee etc. It is important to have adequate fluid to avoid headaches.

Day 5 Day of Colonoscopy

Please follow the instructions that your will receive with your appointment information.

You must take the 3rd litre of **GLYCOPREP-C** on the day of your procedure. You will be instructed **WHAT TIME** to take your final litre of GLYCOPREP-C on your appointment instructions that you will receive from the Endoscopy Unit.

Please follow these instructions carefully.

Some Helpful Hints

- Add diet lemon cordial to the GLYCOPREP-C
- Drink the solution through a straw, suck the occasional barley sugar or butterscotch between drinks
- Apply protective cream, eg. Lanoline around your anal area to reduce excoriation
- Keep warm
- If you feel nauseated whilst taking the preparation stop for ½ hour. Have a hot drink & walk around. Resume drinking & continue as before.
- If you feel bloated, try walking around, suck some peppermints / drink peppermint tea
- If you become ill & have severe pain seek medical advice from your GP or nearest Emergency Medical Centre
- If you have haemorrhoids (piles) use your haemorrhoid cream to reduce discomfort
- Please note that a very clean bowel can only be achieved by drinking all the preparation

If you feel your bowel motion is not clear enough please contact The Prince Charles Hospital Endoscopy Unit on \$\alpha\$3139 4199 after 7.30AM