



Metro North Hospital and Health Service *Putting people first*

Royal Brisbane and Women's Hospital

# Oncologic Emergencies

Matthew Burge

Staff Specialist Cancer Care Services RBWH

July 2017

# Oncologic Emergencies

- Case based talk
- Febrile neutropenia
- Calcium disorders
- Spinal cord compression
- Brain metastases

# Case 1

- 70yr old male
- Diagnosed with metastatic prostate cancer 2015
  - PSA: 300 u/l
  - Bone metastases
  - Gleason score 9 adenocarcinoma
- Commenced on androgen deprivation but developed progression 2016 with new hip pain and rising PSA


# Case 1

- Denosumab (xgeva) commenced  
–120 mg every 4 weeks
- Right hip received radiotherapy
- Docetaxel chemotherapy commenced January 2015
- PSA fell

# Case 1

- Admitted 8 days after 4<sup>th</sup> docetaxel cycle
- Febrile to 39 degrees
- Acute abdominal pain
- Blood pressure 80/40
- Neutrophils 0.46 (2-11)
- Corrected calcium 1.27 (2.1-2.6); ionized 0.78 (1.3-1.5)
- CT: perforated diverticulitis

R  
1  
9  
3



L  
1  
8  
7

# Case 1

- Urgent sigmoid colectomy and Hartmanns
- ICU:
  - Inotropes
  - Calcium chloride IV
  - Broad spectrum IV antibiotics
    - Metronidazole, gentamicin, ampicillin
  - G-CSF
- Gradually recovered
- Calcium took 4 months to return to normal

# Case 1

- Chemotherapy (docetaxel) and denosumab ceased!
- Enzalutamide commenced
- Patients remains quite well to date
- No recurrence of hypocalcaemia



# Febrile Neutropenia

- Very common chemotherapy side effect
- Neutropenia usually 7-14 days after chemotherapy administration
- Various definitions:
  - Fever  $>38$  degrees
  - Neutrophils  $<1.0$
- Urgent assessment required:
  - Attempt to risk stratify
  - Prior episodes.
  - Associated symptoms
    - Mucositis; diarrhoea; pain; confusion; dyspnoea
  - Co-morbidities
  - Antibiotic allergies; recent antibiotics

# Febrile neutropenia

- Examination:
  - Vital signs; oxygenation
  - Infection source
    - Mouth; skin; perineum; central lines; stents
  - Avoid invasive procedures including PR exam
- Treatment:
  - Cultures; radiology as appropriate
  - Urgent intravenous antibiotics

# Calcium disorders

- Hypercalcaemia of malignancy:
  - Squamous cell cancers; Breast; lung; myeloma;
  - Associated with poor prognosis; advanced disease
    - Survival usually measured in months
- Ectopic (tumour cell) secretion of parathyroid related protein most common cause (80%)
  - Don't have to have bone metastases
- Commonly asymptomatic
  - Symptoms can be very non-specific
  - Polyuria; thirst; confusion


# Calcium disorders

- Hypercalcaemia treatment:
  - Re-hydration
  - Bisphosphonates
    - Zoledronic acid 4mg.
- Hypocalcaemia:
  - In oncology, usually drug related.
  - Consider stopping offending agent
  - **Check for hypomagnasemia**
  - Calcium IV replacement guided by symptoms/prolonged QT interval on ECG

# Case 2

- 73 year old male
- Metastatic adenocarcinoma of the lung diagnosed in 2014.
- Treated with various chemotherapy agents; investigational cancer stem cell inhibitor and immune activating antibody.
- Mid 2016:
  - Several weeks of gradually worsening upper back pain
  - Radiating around anteriorly
  - Paraesthesia over left side of trunk/lower limb
  - No weakness or autonomic dysfunction


R  
2  
2  
0



SRA



A  
L



# Case 2

- Dexamethasone 8mg BD commenced immediately
- Urgent radiotherapy to the thoracic spine
- Pain substantially reduced over subsequent weeks
- Dexamethasone gradually weaned.
- Remains ambulatory, being worked up for another clinical trial!



# Spinal cord compression

- Need to consider it as a possibility
  - Anyone with known metastatic malignancy
  - But it might be the first presentation!
  
- Common primary sites:
  - Lung
  - Breast
  - Multiple myeloma
  - Prostate
  
- Thoracic spine > lumbar > cervical
  - From vertebral bone metastases

# Spinal cord compression-symptoms

- Back pain
  - Usually precedes neurologic symptoms
  - Worse lying flat; radicular features
- Neurologic symptoms
  - Weakness
  - Sensory level
  - Urinary retention-late sign
  - ataxia

# Spinal Cord Compression

- MRI is the optimal investigation
  - Entire spine
- Immediate commencement of corticosteroids
  - Optimal dexamethasone dose uncertain
    - 16mg in divided doses
- Surgical decompression
- Radiotherapy
- Chemotherapy
  - Combination of all these


# Spinal cord compression

- **Neurologic status at diagnosis/start of treatment dictates patient outcome.**
- **Starting treatment once paralysed likely futile and will not improve neurologic function.**
- **Early diagnosis and intervention is key.**

# Case 3

- 46 year old lady
- Breast cancer diagnosed 2015.
  - Mastectomy
    - Heavy nodal burden in axilla
  - Adjuvant chemotherapy and radiation to chest wall
- April 2017:
  - 3 day history of severe headache
    - No associated neurological symptoms

R  
I  
A



# Case 3

- Isolated right parietal metastasis causing ventricular effacement
- Dexamethasone 4mg QID commenced with rapid improvement
- Urgent neurosurgical resection
- Followed by whole brain radiotherapy
- Staging CT also demonstrated hepatic metastases
  - Commenced chemotherapy and is doing well

# Brain metastases

- Primary sites:
  - Lung, breast, kidney, melanoma, colorectal
- Increasing incidence:
  - Patients living longer due to better treatments
- Consider prognosis before determining best treatment
  - How much cancer is outside the brain and how treatable is it?
- Corticosteroids; surgery; radiotherapy; systemic therapy
  - Prophylactic anti-epileptics generally not recommended



**Thank you for your attention!**